OPEN ACCESS

University of BRISTOL

Wylde, V., Sayers, A., Odutola, A., Gooberman-Hill, R., Dieppe, PA., & Blom, A. (2017). Central sensitisation as a determinant of patients' benefit from total hip and knee replacement. *European Journal of Pain*, *21*(2), 357–365. https://doi.org/10.1002/ejp.929

Publisher's PDF, also known as Version of record

License (if available):
CC BY-NC-ND

Link to published version (if available):
10.1002/ejp.929

Link to publication record in Explore Bristol Research
PDF-document

This is the final published version of the article (version of record). It first appeared online via Wiley at http://onlinelibrary.wiley.com/doi/10.1002/ejp.929/full?platform=hootsuite. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research
General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/

Supplementary Table 2: Descriptive statistics for patients undergoing total hip replacement (n=90) and total knee replacement (n=75) not included in the analysis because of missing data

	Time	Measure	Mean	(SD)	IQR (25, 50, 75)		
Hips	Pre	PPT mean*	193.93	(90.77)	(121.17,	180.5,	247.17)
		PPT SD#	29.07	(18.85)	(15.45,	25.47,	39.88)
		BMI	29.04	(5.30)	(25.30,	29.03,	32.82)
		Age	65.67	(12.36)	(58,	66,	76)
		WOMAC Pain score	38.95	(18.50)	(25,	40	50)
	Post	WOMAC Pain score	82.25	(21.57)	(75,	90,	100)
Knees	Pre	PPT mean*	203	(99.23)	(131.33,	187.17,	248.33)
		PPT SD#	37.09	(24.87)	(18.52,	29.93,	50.9)
		BMI	32.80	(7.15)	(27.70,	30.88,	36.42)
		Age	69.75	(8.62)	(64,	69,	75)
		WOMAC Pain score	41.47	(16.86)	(25,	40,	55)
	Post	WOMAC Pain score	76.81	(19.65)	(60,	80,	97.5)

*Mean average pressure pain thresholds (PPT) across the 3 replicates

#Mean average PPT standard deviation across the 3 replicates